

The Spectrum of IgM-Associated Peripheral Neuropathies – A Cohort Study from the Neurohaematology Clinic at Queen Square and UCLH

Oliver Tomkins¹, Nicole Japzon¹, Jahanzaib Khawaja¹,

Jindriska Lindsay¹, Charalampia Kyriakou¹, Aisling Carr², Michael P Lunn², Shirley D'Sa¹

1. UCLH Centre for WM and Related Conditions, University College London Hospitals NHS Foundation Trust, London, United Kingdom
2. Centre for Neuromuscular Diseases, National Hospital for Neurology and Neurosurgery, London, United Kingdom

Background

The IgM-associated peripheral neuropathies (IgM-PN) are a heterogeneous collection of disorders, with a causal association with IgM monoclonal gammopathy of undetermined significance (MGUS) or Waldenström's Macroglobulinaemia (WM) via different mechanisms. We describe the distribution, clinical features and treatment outcomes of IgM-PN at our specialist centre.

Methods

A retrospective cohort analysis was undertaken using the UCLH WM patient registry. All patients with a confirmed diagnosis of IgM-PN as deemed by expert assessment, following examination, serological testing and neurophysiological assessment, were included.

Results

A total of 667 patients with IgM gammopathy were reviewed, 98 with IgM MGUS and 569 with WM. IgM-PN was diagnosed in 125 (18.7%) patients, of whom 112 had adequate data available. Baseline clinicopathological details are given in table 1. Patients were diagnosed between years 1997 and 2024, with median follow up of 83 months (3-835m). The underlying disorder was IgM MGUS in 56 (50%), WM in 53 (47.3%), CLL 2 (1.6%) and IgM POEMS 1 (0.8%). No systemic disease was detected in 14/112 cases; isolated flow cytometric disease in 5, and isolated MYD88^{L265P} PCR in 3 cases and 1 case of isolated CXCR4^{WHIM} likely due to a non-L265P MYD88 mutation. Nerve biopsy was done in 22/112 (19.6%) patients.

Frontline treatment (1L) was given in 84/112 (75%), and an additional 2/4 patients with MMN-CB* received intravenous immunoglobulin (IVIg). Median lines of treatment received was 1 (range 1-5). Frontline regimen was rituximab monotherapy (R) in 46, dexamethasone-rituximab-cyclophosphamide (DRC) 22, R-CHOP/CVP in 5, R-bendamustine 3, R-chlorambucil 3, Bortezomib-dexamethasone 2, ibrutinib 1, R-methotrexate-cytarabine 1 (BNS), and lenalidomide-dexamethasone 1 (POEMS).

Progression despite 1L therapy occurred in 19/70 (27.1%) assessable patients, stabilisation in 25/70 (35.7%), and improvement in 26/70 (37.1%). Second-line therapy has been given in 34 cases, including 14 cases of R-retreatment following an initial period of response to 1L therapy. Bruton's Tyrosine Kinase inhibitors (BTKis) were employed in 12 patients, including ibrutinib (n=7), zanubrutinib (n=4) and pirtobrutinib (n=1).

Of patients with anti-MAG neuropathy, 61/80 (76.3%) required therapy, including frontline R in 37 and DRC in 17 cases. Improvement occurred in 22 (41.5%), stabilisation in 20 (37.7%) and progression in 11/53 (20.8%) assessable cases.

On univariate analysis, progressive disease following frontline therapy was significantly more common in patients who do not attain a partial response (p 0.05), AL amyloid (p 0.04) and PN-BNS (p 0.001); AL amyloid remained significant (p 0.039) on multivariate analysis.

Up to date survival information was available in 95 patients; 16/95 (16.8%) are deceased. Median OS for anti-MAG PN 835m (225-NR), with 3- and 5-year OS 100% and 94.8% (89-100%); AL amyloid 162m (144-NR), 3- and 5-year OS 100%.

Conclusion

The incidence of IgM-PN in our cohort was 18.7%, with anti-MAG representing most cases (71%). Clonal burden in the bone marrow is typically small and several patients have disease only on immunophenotyping (n=5) or PCR (n=4). A significant majority of IgM-PN cases is associated with a MYD88^{L265P} mutation. Approximately 73% of patients experienced stabilisation or improvement in PN with frontline therapy.

Table 1 – Clinicopathological Characteristics of Patients with IgM-PN	
Demographics	n = 112
Age, median (range)	66 (30-86)
Male	82 (73.2%)
Underlying Disorder	n = 112
IgM MGUS	56 (50%)
Waldenstrom's Macroglobulinaemia	53 (47.3%)
CLL	2 (1.6%)
IgM POEMS	1 (0.8%)
Pathology	
Paraprotein concentration, median (range) (g/L)	5 (too faint to quantify – 55)
Kappa	98 (87.5%)
Lambda	14 (12.5%)
MYD88 ^{L265P}	36/47 (76.6%)
MYD88 not detected	11/47 (23.4%)
IgM-PN Diagnosis	n = 112
Anti-MAG	80 (71.4%)
AL amyloid	8 (7.1%)
Multifocal motor neuropathy with conduction block* (MMN-CB)	4
Cryoglobulinaemic vasculitis	3
PN Bing-Neel syndrome (BNS)	2
Light chain deposition disease	1
Neurolymphomatosis, high grade transformed	1
CANOMAD	1
IgM POEMS	1
Other, putative IgM-associated	11